



Eur päisch s Patentamt
Eur p an Pat nt Offic
Office européen des brevets



Publication number: **0 635 576 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **94111525.5**

(51) Int. Cl.⁶: **C12P 25/00, C12N 1/20**

(22) Date of filing: **22.07.94**

(30) Priority: **22.07.93 JP 181615/93**

(43) Date of publication of application:
25.01.95 Bulletin 95/04

(84) Designated Contracting States:
CH DE FR GB LI

(71) Applicant: **NIPPON OIL Co. Ltd.**
3-12, Nishi-Shimbashi 1-chome
Minato-ku
Tokyo (JP)

(72) Inventor: **Tsubokura, Akira**
2-228, Kosugi-cho,
Nakahara-ku
Kawasaki-shi,
Kanagawa (JP)
Inventor: **Yoneda, Hisashi**
c/o Nisseki-Ohsato-ryo,
155-72, Honmokuhsato-cho
Naka-ku,
Yokohama-shi,
Kanagawa (JP)
Inventor: **Takaki, Mikihiko, c/o**
Nissekikagaku-Kamihirama-ryo
612-2, Kamihirama
Nakahara-ku,
Kawasaki-shi,
Kanagawa (JP)
Inventor: **Kiyota, Takashi**
94-9, Ichisawa-cho,
Asahi-ku
Yokohama-shi,
Kanagawa (JP)

(74) Representative: **Cohausz & Florack**
Patentanwälte
Kanzlerstrasse 8a
D-40472 Düsseldorf (DE)

EP 0 635 576 A1

(54) **Bacteria belonging to new genus and process for production of carotenoids using same.**

(57) Bacterium belonging to a new genus, and a process for production of carotenoid pigment selected from the group consisting of astaxanthin, adonixanthin, β -carotene, echinenone, canthaxanthin and zeaxanthin, comprising the steps of culturing a bacterium capable of producing at least one of the carotenoid pigments, and recovering an individual carotenoid pigments.

The producer bacterium belongs to new genus.

According to the present process various carotenoids can be produced in an industrial scale. By the present process (3S, 3'S)-astaxanthin can be produced in an almost 100% purity.

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 The present invention provides bacteria belonging to a new genus, and a process for production of carotenoids using a bacterium belonging to the new genus. The carotenoids of the present inventions are natural pigments useful for feed additives, food additives etc. Especially, astaxanthin is valuable in an industrial point of view as feed additives such as a color improver for bred fishes such as salmon, trout, red sea bream etc., and safe natural food additives. In addition, adonixanthin is, if its industrial production
10 process is established, promising as food additives and feed additives as the astaxanthin is.

Further, β -carotene has been used as food additives, feed additives, pharmaceuticals etc., echinenone is promised as food additives, feed additives etc.; canthaxanthin has been used food additives, feed additives, cosmetics etc.; and zeaxanthin has been used as food additives, feed additives etc.

15 2. Related Art

It is known that astaxanthin is contained in fishes such as red sea bream, salmon, trout, etc., and Crustacea such as shrimps, crabs, crawfishes, krills (Carotenoids of Marine Organisms; edt. Nippon Suisan Gakukai, 1978). As microorganisms which produce astaxanthin, red yeast *Phaffia rhodozyma* -
20 (Phytochemistry, 15, 1009, 1976), *Brevibacterium* (Journal of General and Applied Microbiology, 15, 127, 1969), and green alga *Haematococcus pluvialis* (Phytochemistry, 20, 2561, 1981) are known. As chemical synthetic processes, conversion of β -carotene (Pure Appl. Chem. 57, 741, 1985) and synthesis from phosphonium salt (Helv. Chim. Acta. 64, 2436, 1981) are known.

However, the known process for production of astaxanthin are not advantageous due to high cost,
25 because a content of astaxanthin in natural products such as krills, crawfishes, etc. is very low, and extraction thereof is difficult. In addition, stable availability of the resources is problematic. In addition, growth rate of red yeast, *Phaffia rhodozyma*, is low, productivity of astaxanthin by the red yeast is low, and extraction of astaxanthin is difficult since this yeast has hard cell wall. Therefore, industrial production of astaxanthin from the red yeast is difficult.

30 A growth rate of green alga, *Haematococcus pluvialis*, is also low, the culture is easily contaminated, and extraction of astaxanthin is difficult. Therefore industrial production of astaxanthin from the red alga is difficult.

It is known that adonixanthin is contained in goldfishes and carps (Carotenoids of Marine Organisms, Nippon Suisan Gakukai, 1978), but it is believed that chemical synthesis of adonixanthin is difficult. No
35 industrial process for production of adonixanthin is known.

As production processes for β -carotene, although synthesis from β -ionone (Pure & Appl. Chem. 63(1), 45, 1991), and extraction from green or yellow vegetables such as carrot, sweet potato, pumpkin etc. are known (Natural Coloring Agent Handbook, Korin (1979), edt. by editorial committee of Natural Coloring Agent Handbook), production cost of these processes is high. As processes for production of β -carotene by
40 microorganisms, the production by an alga *Dunaliella* (J. Appl. Bacteriol., 70, 181, 1991), and the production by a fungus *Blakeslea* (J. Appl. Bacteriol., 70, 181, 1991) are known. However, production of β -carotene by bacteria is not known.

Echinenone is extracted from natural products, for example, starfishes such as crown-of-thorns starfish, the internal organs of fishes such as red sea bream, sea urchin, the internal organs of Crustacea such as
45 lobster, etc. (Carotenoids of Marine Organisms, edt. Nippon Suisan Gakukai, 1987). However, the production of echinenone by microorganisms is not known.

Canthaxanthin is known to be contained in a sort of mushrooms (Botanical Gazette, 112, 228 - 232, 1950), fishes, Crustacea etc. (Carotenoids of Marine Organisms, edt. Nippon Suisan Gakukai, 1978). The production of echinenone by microorganisms are exemplified by the production by microorganisms
50 belonging to the genus *Brevibacterium* (Applied and Environmental Microbiology, 55(10), 2505, 1989), and by microorganisms belonging to the genus *Rhodococcus* (Japanese Unexamined Patent Publication No. 2-138996). In addition, as chemical synthetic processes, oxidation of β -carotene (J. Amer. Chem. Soc., 78, 1427, 1956) and synthesis from novel compound 3-oxo-C₁₅ phosphonium salt (Pure & Appl. Chem. 51, 875, 1979) are known.

55 As processes for production of zeaxanthin, a chemical synthesis starting from a hydroxy ketone obtained by asymmetric reduction of oxoisophorone (Pure & Appl. Chem., 63(1), 45, 1991), extraction from corn seeds (Seitai Shikiso, 1974, Asakura Shoten), and a process using *Flavobacterium* (Carotenoids, In Microbial Technology, 2nd edn. Vol. 1, 529 - 544, New York: Academic Press) are known.

SUMMARY OF INVENTION

The present invention provides bacteria belonging to a new genus.

The present inventors found that novel bacterial strains not belonging to any known genera produce
5 various kinds of carotenoid pigments.

Accordingly, the present invention further provide a process for production of a carotenoid selected from the group consisting of astaxanthin, adonixanthin, β -carotene, echinenone, canthaxanthin and zeaxanthin comprising culturing a microorganism belonging to a new genus and capable of producing at least one of said carotenoids, and recovering a carotenoid or a combination of the carotenoids from the culture.

10

BRIEF EXPLANATION OF DRAWINGS

Figure 1 shows a ^{13}C nuclear magnetic resonance spectrum of astaxanthin produced by the present process.

15 Fig. 2 shows a mass spectrum of astaxanthin produced by the present process.

Fig. 3 shows a ^{13}C nuclear magnetic resonance of adonixanthin produced by the present process.

Fig. 4 shows a mass spectrum of adonixanthin produced by the present process.


Fig. 5 shows a profile of elution of carotenoid pigments of the present invention in a high performance liquid chromatography.

20 Fig. 6 shows a phylogenetic tree of E-396 strain and close species.

DETAILED DESCRIPTION OF INVENTION

The carotenoids produced according to the present invention are represented by the following formula:

25



30

35

Astaxanthin


{3,3'-Dihydroxy- β,β -Carotene-4,4'-dione}

40


45

50


55




Adonixanthin
(3,3'-Dihydroxy- β,β -Caroten-4-one)




β -Carotene
(β,β -Carotene)



Echinenone
(β,β -Caroten-4-one)



Canthaxanthin
(β,β -Carotene-4,4'-dione)



Zeaxanthin

((3R,3'R)- β,β -Carotene-3,3'-diol)

According to the present invention any bacteria capable of producing the above-mentioned carotenoids and having the following properties can be used. A group of bacteria having such properties do not belong to any of known genera described in Bergey's Manual of Systematic Bacteriology, and therefore is a group of bacteria belonging to a new genus.

(1)	Morphology	polymorphic rod
(2)	Motility	motile
(3)	Flagella	peritricous
(4)	Spore formation	none
(5)	Gram stain	negative
(6)	Production of pigment	positive (water insoluble)
(7)	Oxidase	positive
(8)	Catalase	positive
(9)	Behavior toward oxygen	aerobic
(10)	Fermentation of glucose	negative
(11)	Production of 3-ketolactose	negative
(12)	Quinone type	Q-10
(13)	GC content of intracellular DNA	64 - 69 molar%
(14)	Formation of slime	
	Glucose	negative
	Sucrose	negative
(15)	Presence of sphingolipid	negative
(16)	Presence of bacteriochlorophyll	negative

Among those bacteria, as a particular microorganism E-396 strain can be mentioned. This strain was newly isolated by the present inventors, and has been deposited with National Institute of Bioscience and Human-Technology Agency of Industrial Science and Technology, 1-3 Higashi 1-chome, Tsukuba-shi, Ibaraki-ken Japan, on April 27, 1993 as FERM BP-4283 under the Budapest Treaty.

This strain has the following properties:

(a) Morphological characters

(1)	Shape and size of cells rod, 0.5 - 0.75 \times 1.0 - 5.0 μ m	
(2)	Polymorphism	present
(3)	Motility	present, peritricous
(4)	Spores	none

(b) Cultural characters

(1) Bouillon agar plate culture

Circular orange glossy colony is formed. Diffusible pigment is not formed.

(2) Bouillon liquid culture

Growth is somewhat poor; totally cloudy; precipitated. Surface growth is not present.

EP 0 635 576 A1

(3) Bouillon gelatin stab culture
Growth is poor; gelatin is not liquefied.

(4) Litmus milk
Not changed

5 (c) Physiological characteristics

	(1) Gram stain	negative
	(2) Reduction of nitrate	negative
10	(3) Denitrification	negative
	(4) MR test	negative
	(5) VP test	negative
15	(6) Production of indole	negative
	(7) Production of hydrogen sulfide	negative

20

25

30

35

40

45

50

55

EP 0 635 576 A1

	(8)	Hydrolysis of starch	negative
	(9)	Utilization of citric acid	
5		Koser medium	negative
		Christensen medium	negative
	(10)	Utilization of inorganic nitrogen sources	
		Nitrate	negative
10		Ammonium salt	positive
	(11)	Production of pigment	positive
		(water insoluble)	
15	(12)	Urease	negative
	(13)	Oxidase	positive
	(14)	Catalase	positive
20	(15)	Range of growth	
	(i)	pH	growing at pH 6 to 9 (optimum pH 7)
25		(ii) Temperature	growing at 10°C to 33°C (optimum 28°C)
30	(16)	Behavior toward oxygen	aerobic
	(17)	O-F test	oxidation
35			(Hugh- Leifson medium supplemented with 0.1% yeast extract)
40			
45	(18)	Formation of acid or gas from saccharides (Hugh-Leifson medium supplemented with 0.1% yeast extract)	
		Acid	Gas
50	(1)	L-Arabinose	+
	(2)	D-Xylose	+

55

EP 0 635 576 A1

	(3)	D-Glucose	+	-
	(4)	D-Mannose	+	-
5	(5)	D-Fructose	-	-
	(6)	D-Galactose	+	-
	(7)	Maltose	+	-
	(8)	Sucrose	+	-
10	(9)	Lactose	+	-
	(10)	Trehalose	+	-
	(11)	D-Sorbitol	+	-
15	(12)	D-Mannitol	+	-
	(13)	Inositol	-	-
	(14)	Glycerin	+	-
20	(15)	Starch	-	-

(d) Other physiological characteristics

25	(1)	Decomposition of esculin	positive
	(2)	Arginine dihydrolase	negative
	(3)	Formation of 3-ketolactose	negative
30	(4)	PNPG	positive
	(5)	Formation of slime	
		Glucose	negative
		Sucrose	negative
35	(6)	Utilization (by API 20NE kit)	
	(1)	Glucose	positive
	(2)	L-Arabinose	negative
40	(3)	D-Mannose	positive
	(4)	Mannitol	positive
	(5)	N-Acetyl-D-glucosamine	negative
45	(6)	Maltose	positive
	(7)	Potassium gluconate	positive
	(8)	n-Capronate	negative
	(9)	Adipic acid	negative
50	(10)	dl-Malic acid	positive
	(11)	Sodium citrate	negative
55	(12)	Phenyl acetate	negative

e) Chemotaxonomic characters

- (1) Intracellular GC content of 67 molar%
DNA (HPLC method)

- (2) Quinone type Q-10
(3) Bacteriochlorophyll
Production in anaerobic condition
non growth
Production in aerobic condition
negative
(4) Presence of sphingolipid negative
(5) Cellular fatty acid composition

Fatty acid	Ratio (%)
C16:0	0.8
C16:1	0.1
C17:0	0.5
C18:0	9.0
C18:1	81.4
C19:0	0.4
2-OH C14:0	0.2
3-OH C14:0	0.6

- (6) DNA-DNA homology with analogous species
(HPLC method)

Strains tested	Homology (%)
Flavobacterium okeanokoites IFO 12536	2
Flavobacterium resinovorum ATCC 33545	23
Flavobacterium aquatile IFO 15052	0
Agrobacterium tumefaciens IFO 15193	0
Agrobacterium radiobacter IFO 12532	7
Mycoplana bullata IFO 13290	32
Mycoplana segnis IFO 15250	26
Mycoplana ramosa IFO 15249	9
Mycoplana dimorpha IFO 13291	11
Xanthobacter flavus IFO 14759	24
Sphingomonas paucimobilis IFO 13935	6

- (7) Nucleotide sequence of DNA coding for 16S
ribosome RNA
Shown in SEQ ID:1

From the above result, the strain E-396 was considered to be a bacterium belonging to the genus Agrobacterium since it is aerobic, gram negative and peritricous rod, but this was negated due to the results of pigment productivity, slime-forming ability and DNA-DNA homology. In addition, the strain was considered to be a bacterium belonging to the genus Sphingomonas or a photosynthesis bacterium from the color of colony, but they were also negated because sphingolipid and bacteriochlorophyll were not detected.

Some stored strains assumed to be close to the strain E-396 in the similarity of colony color, peritricous flagella, GC content and quinone type were selected, and DNA-DNA homology between the strain E-396 and the selected strains was tested. As a result, there is no genus which shows the similarity to the strain E-396 in the DNA-DNA homology. In addition, a phylogenetic tree was prepared on the basis of the nucleotide sequence of 16S ribosome RNA of the strain E-396 according to the neighbor-joining method. As a result, it was found that the strain E-396 is independent from any close genus. Therefore it was confirmed that the strain E-396 is a bacterium belonging to a new genus but not belonging to any known genus.

As further particular microorganism, the strain A-581-1 is mentioned. This strain was newly isolated by the present inventors, and has been deposited with National Institute of Bioscience and Human Technology Agency of Industrial Science and Technology, 1-3 Higashi 1-chome, Tsukuba-shi, Ibaraki-ken Japan, on May 20, 1994 as FERM BP-4671 under the Budapest Treaty.

This strain has the following properties:

(a) Morphological characters

(1)	Shape and size of cell rod, 0.5 - 1.0 × 3.5 μm	
(2)	Polymorphism	present
(3)	Motility	present, peritricous
(4)	Spores	none

(b) Cultural characters

(1) Bouillon agar plate culture Circular orange glossy colony is formed. Diffusible pigment is not formed.

(2) Bouillon liquid culture

Totally cloudy; precipitated. Surface growth is not present.

(3) Bouillon gelatin stab culture

Growth is poor; gelatin is not liquefied.

(4) Litmus milk

Not changed

(c) Physiological characteristics

	(1)	Gram stain	negative
5	(2)	Reduction of nitrate	negative
	(3)	Denitrification	negative
	(4)	MR test	negative
10	(5)	VP test	negative
	(6)	Production of indole	negative
	(7)	Production of hydrogen sulfide	negative
15	(8)	Hydrolysis of starch	negative
	(9)	Utilization of citric acid	
		Koser medium	negative
20		Christensen medium	negative
	(10)	Utilization of inorganic nitrogen sources	
		Nitrate	negative
25		Ammonium salt	positive
	(11)	Production of pigment	positive (water insoluble)
30	(12)	Urease	negative
	(13)	Oxidase	positive
	(14)	Catalase	positive
	(15)	Range of growth	
35		(i) pH	growing at pH 6 to 10 (optimum pH 8)
40		(ii) Temperature	growing at 10°C to 33°C (optimum 28°C)

- (16) Behavior toward oxygen aerobic
- (17) O-F test oxidation (Hugh-Leifson medium supplemented with 0.1% yeast extract)
- (18) Formation of acid or gas from saccharides (Hugh-Leifson medium supplemented with 0.1% yeast extract)

		Acid	Gas
(1)	L-Arabinose	+	-
(2)	D-Xylose	+	-
(3)	D-Glucose	+	-
(4)	D-Mannose	+	-
(5)	D-Fructose	+	-
(6)	D-Galactose	+	-
(7)	Maltose	+	-
(8)	Sucrose	+	-
(9)	Lactose	+	-
(10)	Trehalose	+	-
(11)	D-Sorbitol	-	-
(12)	D-Mannitol	+	-
(13)	Inositol	-	-
(14)	Glycerin	+	-
(15)	Starch	-	-

(d) Other physiological characteristics

5	(1)	Decomposition of esculin	positive
	(2)	Arginine dihydrolase	negative
	(3)	Formation of 3-ketolactose	negative
	(4)	PNPG	positive
10	(5)	Formation of slime	
		Glucose	negative
		Sucrose	negative
15	(6)	Utilization (by API 20NE kit)	
	(1)	Glucose	positive
	(2)	L-Arabinose	positive
	(3)	D-Mannose	negative
20			
	(4)	Mannitol	positive
	(5)	N-Acetyl-D-glucosamine	negative
25	(6)	Maltose	negative
	(7)	Potassium gluconate	negative
	(8)	n-Capronate	negative
30	(9)	Adipic acid	negative
	(10)	dl-Malic acid	positive
	(11)	Sodium citrate	negative
35	(12)	Phenyl acetate	negative

(e) Chemotaxonomic characters

40	(1)	Intracellular GC content of DNA (HPLC method)	66 molar%
	(2)	Quinone type	Q-10
	(3)	Bacteriochlorophyll	
		Production in anaerobic condition	non growth
		Production in aerobic condition	negative
45	(4)	Presence of sphingolipid	negative
	(5)	DNA-DNA homology with the strain E-396 (HPLC method)	homology 56%

From the above results as well as the facts that the strains A-581-1 has the following characteristics as the strain E-396 has, and that the strain A-581-1 has high DNA-DNA homology with the strain E-396, it is determined that the strain A-581-1 belongs to the new genus to which the strain E-396 also belongs.

(1)	Morphology	polymorphic rod
(2)	Motility	motile
(3)	Flagella	peritricous
(4)	Spore formation	none
(5)	Gram stain	negative
(6)	Production of pigment	positive (water insoluble)
(7)	Oxidase	positive
(8)	Catalase	positive
(9)	Behavior toward oxygen	aerobic
(10)	Fermentation of glucose	negative
(11)	Production of 3-ketolactose	negative
(12)	Quinone type	Q-10
(13)	GC content of intracellular DNA	64 - 69 molar%
(14)	Formation of slime	
	Glucose	negative
	Sucrose	negative
(15)	Presence of sphingolipid	negative
(16)	Presence of bacteriochlorophyll	negative

Medium for production of carotenoids using the present microorganisms is, for example, as follow. Namely, it contains a carbon source, a nitrogen source and inorganic salts necessary for the growth of producer microorganisms, as well as if necessary special required substances (for example, vitamins, amino acids, nucleic acids etc.). As the carbon sources, sugars such as glucose, sucrose, lactose, fructose, trehalose, mannose, mannitol, maltose, etc., organic acids such as acetic acid, fumaric acid, citric acid, propionic acid, malic acid, pyruvic acid, malonic acid; alcohols such as ethanol, propanol, butanol, pentanol, hexanol, isobutanol, glycerol; oil or fat such as soybean oil, rice bran oil, olive oil, corn oil, sesame oil, linseed oil, and the like are mentioned. Amount of the carbon source added varies according to the kind of the carbon source, and usually 1 to 100g, preferably 2 to 50g per 1 l medium.

As the nitrogen sources, for example, potassium nitrate, ammonium nitrate, ammonium chloride, ammonium sulfate, ammonium phosphate, ammonia, urea etc. are used alone or in combination. Amount of the nitrogen source added varies according to the kind of the nitrogen source, and usually 0.1 to 30g, and preferably 1 to 10g per 1 l medium.

As the inorganic salts, potassium dihydrogen phosphate, dipotassium hydrogen phosphate, disodium hydrogen phosphate, magnesium sulfate, magnesium chloride, ferric sulfate, ferrous sulfate, ferric chloride, ferrous chloride, manganous sulfate, manganous chloride, zinc sulfate, zinc chloride, cupric sulfate, calcium chloride, calcium carbonate, sodium carbonate etc. may be used alone or in combination. Amount of inorganic acid varies according to the kind of the inorganic salt, and usually 0.001 to 10g per 1 l medium.

As special required substances, vitamins, nucleic acids, yeast extract, peptone, meat extract, malt extract, corn steep liquor, soybean meal, dried yeast etc. may be used alone or in combination.

Amount of the special required substance used varies according to the kind of the substance, and usually 0.2g to 200g, and preferably 3 to 100g per 1 l medium. A pH value of a medium is adjusted to pH 2 to 12, preferably 6 to 9. Culturing is carried out at temperature of 15 to 80 °C, and preferably 20 to 35 °C, usually for 1 to 20 days, and preferably 2 to 8 days, under an aerobic condition provided by shaking or aeration/agitation.

Finally the present compound may be isolated and purified from the culture. Namely, microbial cells are separated from the culture by a conventional means such as centrifugation or filtration, and the cells are subjected to an extraction with a solvent. Since a small amount of carotenoids is dissolved in a supernatant or filtrate, the carotenoids also may be recovered from them. As a solvent for the extraction, any substance in which the present compound is soluble can be used. For example, organic solvents such as acetone, chloroform, dichloromethane, hexane, cyclohexane, methanol, ethanol, isopropanol, benzene, carbon disulfide, diethyl ether etc. are used, and preferably chloroform, dichloromethane, acetone, methanol, ethanol or isopropanol is used. The purification can be carried out by conventional procedures such as absorption, elution, dissolving and the like, alone or preferably in combination.

According to the present invention, in many cases, astaxanthin, adonixanthin, β -carotene, echinenone, canthaxanthin and zeaxanthin are simultaneously produced and present in a culture product. Accordingly, in an embodiment of the present invention, any one of the above-mentioned carotenoids can be singly

obtained by the above-mentioned procedure. Alternatively, a mixture of the carotenoids also can be obtained. In this way, the process for carotenoid production of the present invention includes a process for production of an individual carotenoid and a process for production of a mixture of the carotenoids.

Astaxanthin and adonixanthin can be separated from each other according to a conventional procedure
 5 for mutual separation of carotenoids, such as adsorption/elution column chromatography, differential extraction, counter current extraction, differential crystallization.

In addition, for production of an individual carotenoid, the desired carotenoid may be preferentially produced by controlling medium composition, culture conditions and the like.

For example, a ratio of carotenoids produced can be changed by changing an aerobic condition. For
 10 example, a ratio of carotenoids produced may be changed by an amount of a medium or a rate of shaking in flask-shaking culture, or by changing a rate of air supply or a rate of agitation in aeration/agitation culture. As a particular example, there is a tendency in flask culture that an amount of astaxanthin produced increases, while an amount of adonixanthin decreases, as a volume of a culture medium in a flask increases.

Alternatively, for preferential production of a particular carotenoid, a producer microorganism can be
 15 improved by a mutation such as artificial mutation of the producer microorganism so that a mutant microorganism preferentially produces the desired carotenoid among others. Such mutation treatments include, for example, physical methods such as X-ray radiation, UV radiation and the like; chemical methods such as the use of N-methyl-N'-nitro-N-nitrosoguanidine (NTG), ethylmethane sulfonate (EMS);
 20 and a biological methods such as gene recombination technique. Processes for production of the carotenoids using such an improved mutant is included in the present process for production of carotenoids.

In astaxanthin produced by the present process as described above, purity of (3S, 3'S)-astaxanthin is almost 100%. It is known that a ratio of (3S, 3'S)-astaxanthin in astaxanthin contained in natural products
 25 such as crawfish, *Haematococcus*, salmon, trout, red sea bream is high. On the other hand, it is known that *Phaffia rhodozyma* contains (3R, 3'R)-astaxanthin in a high ratio, which absolute configuration is the opposite of that of astaxanthin contained in most of natural products.

Almost 100% of astaxanthin produced by the present process is (3S, 3'S)-astaxanthin whose absolute configuration is same as that of a majority of naturally occurring astaxanthin, and therefore astaxanthin
 30 produced by the present process is industrially valuable. In addition, although chemical synthesis of (3S, 3'S)-astaxanthin is known (Helv. Chim. Acta, 61, 2609, 1978), since optically pure (4R, 6R)-4-hydroxy-2,2,6-trimethylcyclohexanone is used as a starting material, the process is of high cost, and industrially not advantageous.

In addition, astaxanthin produced by the present process contains all-trans astaxanthin in a high ratio,
 35 with a ratio of all-trans form:cis form being 92:8 to 96:4. The all-trans astaxanthin is of natural form, and the present producer microorganisms are advantageous in that they produce natural type astaxanthin. Where cis-astaxanthin is necessary, this can be obtained from all-trans astaxanthin according to a known process, while it is difficult to prepare all-trans astaxanthin from cis-astaxanthin.

¹³C nuclear magnetic resonance spectrum and mass spectrum of the astaxanthin produced by the
 40 present process are shown in Figs. 1 and 2 respectively. ¹³C nuclear magnetic resonance spectrum and mass spectrum of adonixanthin produced by the present process are shown in Figs. 3 and 4 respectively.

EXAMPLES

45 Next, the present invention is explained by Examples. However the scope of the present invention should not be limited to the Examples.

Example 1

50 One gram of soil obtained in Midori-ku, Kanagawa-prefecture, Japan was suspended in 5 mL of sterilized physiological saline, the supernatant was diluted 100 fold and plated on a buillon agar plate medium, followed by culturing at 30 °C for 3 days. Colonies with orange color were selected so as to obtain a strain E-396 (FERM BP-4283).

From the above-described taxonomical properties, it is confirmed that the strain E-396 (FERM BP-4283)
 55 is a bacterium belonging to a new genus.

Example 2

One gram of soil obtained in Midori-ku, Kanagawa-prefecture, Japan was suspended in 5 mL of sterilized physiological saline, the supernatant was diluted 100 fold and plated on a buillon agar plate medium, followed by culturing at 30 °C for 5 days. Colonies with orange color were selected so as to obtain a strain A-581-1 (FERM BP-4671).

From the above-described taxonomical properties, it is determined that the strain A-581-1 (FERM BP-4671) has the similar properties as the E-396 and belongs to the new genus to which the strain E-396 belongs, because the A-581-1 has high DNA-DNA homology with the E-396.

Example 3

Ten mL of a medium containing 2 g/L glucose, 3 g/L meat extract, 10 g/L pepton and 5 g/L sodium chloride was put into a test tube having a diameter of 18 mm, which was then autoclaved at 121 °C for 15 minutes. A piece of cells of E-396 strain (FERM BP-4283) was inoculated into the medium, and cultured at 30 °C for 6 days at 300 rpm with shaking. 100 tubes of the cultured broth (1L) was centrifuged to obtain cells which were then extracted with 500 mL of acetone, and to the separated acetone layer were added 500 mL of hexane and 500 mL of 0.85% saline, and the whole was mixed. The upper layer is recovered, and the solvent was evaporated off at 35 °C under a reduced pressure.

The extract thus obtained was adsorbed on a silica gel column, and eluted with a mixture of benzene/ethyl acetate/methanol (15:4:1) to obtain an astaxanthin fraction, from which then the solvent was evaporated off under a reduced pressure. The extract was dissolved in a small amount of pyridin, and water was dropwise added to the pyridin solution to crystallize astaxanthin, and 1.2 mg of crystallin astaxanthin was obtained. Astaxanthin thus obtained was identical with authentic astaxanthin in its IR spectrum, mass spectrum, ¹³C nuclear magnetic resonance spectrum and absorption spectrum.

Example 4

Ten mL of a medium having the composition as shown in Table 1 was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes.

Table 1

Composition	Amount added
Yeast extract	20 g/L
Sucrose	10 g/L
KH ₂ PO ₄	1.5 g/L
Na ₂ HPO ₄	1.5 g/L
MgSO ₄ · 7H ₂ O	0.5 g/L
FeSO ₄ · 7H ₂ O	0.01 g/L
CaCl ₂ · 2H ₂ O	0.01 g/L
pH 7.0 (adjusted with Na ₂ CO ₃)	

A piece of cells of E-396 (FERM BP-4283) was inoculated into the tubes and cultured at 30 °C for 2 days with reciprocating shaker at 300 rpm. 10 mL of the culture was inoculated into 100 mL of a medium having the same composition as described above, in a 500 mL Sakaguchi flask, and reciprocating culturing at 100 rpm was carried out at 30 °C for 2 days. Next, 100 mL of the culture thus prepared was inoculated into 3.0 L of a medium having the same composition as described above in a 5.0 L fermentor, and culturing was aerobically carried out at 30 °C and 1.0 vvm for 52 hours.

Next, 3 L of the culture was inoculated into 35 L of a medium having the same composition as described above (except that 10 g/L yeast extract, 20 g/L sucrose and 2.5 g/L NH₄NO₃ were used) in a 50 L fermentor, and culturing was aerobically carried out at 30 °C, 250 rpm and 1.0 vvm for 18 hours. 33 L of the cultured broth thus obtained was centrifuged to obtain 790g of wet cells, which were then washed with 1.3 L of methanol and extracted three times with 0.8 L of chloroform. Astaxanthin was purified from the pigment extract according to the same procedure as described in Example 3 to obtain 10 mg of crystalline astaxanthin.

EP 0 635 576 A1

The astaxanthin thus obtained conformed to an authentic astaxanthin in its IR spectrum, mass spectrum ¹³C nuclear magnetic resonance spectrum, and absorption spectrum. In addition, the absolute configuration of astaxanthin thus obtained was determined by known method (J. High Resolut. Chromatogr. Chromatogr Commun., 2, 195, 1979). As a result, 100% of astaxanthin was (3S, 3'S)-astaxanthin, and a ratio of all-trans-astaxanthin and cis-astaxanthin was 95:5. For the purpose of comparison, a result of analysis for astaxanthin produced by other processes was shown in Table 2.

Table 2

	3R, 3'R form	meso form	3S, 3'S form
Product by E-396	0.0%	0.0%	100.0%
Carophyll pink, Roche	24.8%	50.2%	25.0%
Extract from crawfish	17.6%	28.4%	53.9%

Example 5

Ten mL of a medium having the same composition as shown in Table 1 (except that 30 g/L sucrose was used) was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes. A piece of cells of E-396 (FERM BP-4283) was inoculated into the medium and cultured at 25 °C for 5 days with reciprocating at 300 rpm. 10 mL of the culture was centrifuged to obtain cells, which were then extracted with 10 mL of acetone. To the separated acetone layer, were added 10 mL of hexane and 10 mL of 0.85% saline, and the upper layer was recovered and the solvent was evaporated off at 35 °C under a reduced pressure.

The pigment extract thus obtained was analysed for its carotenoid content by a high performance liquid chromatography, and a result shown in Table 3 was obtained. The analytical condition was as follows: A column was prepared by linking two Wakosil 5SIL-120 columns (Wako Pure Chemical) of 4.6 mm I.D × 250 mm, and the mobile phase was a mixture of hexane/dichloromethane/methanol (10:8:1). Detection of carotenoids was carried out by absorption at 470 nm, and quantitative analysis was carried out by comparing an area of the peak of astaxanthin in a test sample and that of an authentic preparation. In addition, the absolute configuration of astaxanthin was determined in accordance with the same procedure as described above. As a result, 100% of astaxanthin produced was (3S, 3'S)-astaxanthin, and a ratio of all-trans form and cis form was 95:5.

Table 3

Astaxanthin mg/L	Adonixanthin mg/L	Total carotenoids mg/L
19.4	13.5	61.9

Example 6

Ten mL of a medium having the same composition as shown in Table 1 (except that 30 g/L sucrose was used) was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes. A piece of cells of E-396 (FERM BP-4283) was inoculated into the medium, and cultured at 25 °C for 2 days with reciprocating at 300 rpm. The culture was inoculated at an amount of 1% by volume into different volumes between 25 mL and 200 mL of media having the same composition as described above in 500 mL conical flasks, and rotary shaking culture was carried out at 25 °C, for 5 days at 120 rpm. Extraction and analysis of carotenoids from the culture broths were carried out according to the same procedure as described in Example 5. Amounts of astaxanthin, adonixanthin and total carotenoids are shown in Table 4.

Table 4

Volume of Medium in conical flask mL	Astaxanthin mg/L	Adonixanthin mg/L	Total carotenoids mg/L
25	5.6	56.2	63.3
50	16.4	31.5	76.2
100	6.8	20.9	53.4
200	11.6	5.8	34.7

Example 7

The pigment extract obtained in Example 4 was adsorbed on a silica gel column, and an adonixanthin fraction was eluted with a mixed solvent of benzene, ethyl acetate and methanol (15:4:1), and the solvent was evaporated off under a reduced pressure. The extract was dissolved in ethanol at 50 °C, and the solution was allowed to stand at 4 °C for one day to crystallize adonixanthin. 190 mg of crystalline adonixanthin was obtained. The adonixanthin thus obtained conformed to authentic adonixanthin in its IR spectrum, mass spectrum, ¹³C NMR spectrum and absorption spectrum.

Example 8

Ten mL of a medium having the same composition as shown in Table 1 (except that 30 g/L sucrose was used) was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes. A piece of cells of E-396 (FERM BP-4283) was inoculated into the medium and cultured at 28 °C for 5 days with shaking. Carotenoids were extracted from the culture according to the same procedure as described in Example 5, and the carotenoids were analyzed by a high performance liquid chromatography. Elution time of each peak and the wave length at the maximum absorption of each peak in the same solvent as elute are shown in Table 5.

Table 5

Elution time and wave length of maximum absorption for each peak		
Peak No.	Elution time (min)	λ_{max} nm
No. 1	6.0	(428) 456 481
No. 2	6.3	464
No. 3	6.8	474
No. 4	10.4	(426) 455 482

Elution time and wave length of maximum absorption were measured for authentic β -carotene and canthaxanthin according to the same procedure as described above. As a result, the measurements of the peak No. 1 and peak No. 3 well conformed to those of β -carotene and canthaxanthin respectively. Therefore, the peak No. 1 and peak No. 3 were identified as those representing β -carotene and canthaxanthin respectively. In addition, other peaks were compared with the maximum absorption peaks described in Davies B.H. 1976, Carotenoids, 38 - 165, In T.W. Goodwin (ed.), Chemistry and Biochemistry of Plant Pigments, Vol. 2, Academic Press, Inc. (London), Ltd., London. As a result, the peak No. 2 and the peak No. 4 were identified as those representing echinenone and zeaxanthin respectively.

Example 9

Ten mL of brain heart infusion bouillon medium (Eiken Chemical Co., Ltd., pH value was adjusted to 10 with Na₂CO₃) was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes. A piece of cells of A-581-1 strain (FERM BP-4671) was inoculated into the medium and cultured at 33 °C for 4 days with shaking. Carotenoids were extracted from the culture, and quantitatively analyzed by high performance liquid chromatography, according to the same procedure as described in Example 5. A result is shown in Table 6.

EP 0 635 576 A1

Table 6

Astaxanthin mg/L	Adonixanthin mg/L	Total carotenoids mg/L
0.30	0.45	0.72

In addition, the absolute configuration of astaxanthin thus obtained was determined according to the same procedure as described in Example 4, and it was confirmed that 100% of astaxanthin produced was (3S, 3'S)-astaxanthin, and that a ratio of all-trans astaxanthin and cis-astaxanthin was 95:5. A result is shown in Table 7.

Table 7

3R, 3'R form	meso form	3S, 3'S form
0.0%	0.0%	100.0%

Example 10

Ten mL of a medium having the composition shown in Table 8 was put into each of test tubes having a diameter of 18 mm, and autoclaved at 121 °C for 15 minutes.

Table 8

Composition	Amount added
Glucose	10 g/L
Polypepton	5 g/L
Yeast extract	5 g/L
KH ₂ PO ₄	1 g/L
MgSO ₄ · 7H ₂ O	0.2 g/L
pH 8 (adjusted with Na ₂ CO ₃)	

A piece of A-581-1 strain (FERM BP-4671) was inoculated into the medium, and cultured at 28 °C for 4 days with shaking. Carotenoids were extracted from the culture and quantitatively analyzed by high performance liquid chromatography according to the same procedure as described in Example 5. A result is shown in Table 9.

Table 9

Astaxanthin mg/L	Adonixanthin mg/L	Total carotenoids mg/L
0.17	0.64	2.81

SEQUENCE LISTINGS

SEQ ID NO.: 1

5 SEQUENCE LENGTH: 1452

SEQUENCE TYPE: nucleic acid

10 STRANDNESS: double

MOLECULE TYPE: DNA coding for rRNA

	AGTTTGATCC	TGGCTCAGAA	CGAACGCTGG	CGGCAGGCTT	AACACATGCA	50
15	AGTCGAGCGA	GACCTTCGGG	TCTAGCGGCG	GACGGGTGAG	TAACGCGTGG	100
	GAACGTGCCC	TTCTCTACGG	AATAGCCCCG	GGAAACTGGG	AGTAATACCG	150
	TATACGCCCT	TTGGGGGAAA	GATTTATCGG	AGAAGGATCG	GCCCGCGTTG	200
20	GATTAGGTAG	TTGGTGGGGT	AATGGCCAC	CAAGCCGACG	ATCCATAGCT	250
	GGTTTGAGAG	GATGATCAGC	CACACTGGGA	CTGAGACACG	GCCCAGACTC	300
	CTACGGGAGG	CAGCAGTGGG	GAATCTTAGA	CAATGGGGGC	AACCCTGATC	350
25	TAGCCATGCC	GCGTGAGTGA	TGAAGGCCTT	AGGGTTGTAA	AGCTCTTTCA	400
	GCTGGGAAGA	TAATGACGGT	ACCAGCAGAA	GAAGCCCCGG	CTAACTCCGT	450
30	GCCAGCAGCC	GCGGTAATAC	GGAGGGGGCT	AGCGTTGTTC	GGAATTACTG	500
	GGCGTAAAGC	GCACGTAGGC	GGAAGTGGAA	GTCAGAGGTG	AAATCCCAGG	550
	GCTCAACCTT	GGAAGTGCCT	TTGAAACTAT	CAGTCTGGAG	TTGAGAGAG	600
35	GTGAGTGGAA	TTCCGAGTGT	AGAGGTGAAA	TTCGTAGATA	TTGAGAGGAA	650
	CACCAAGTGGC	GAAGGCGGCT	CACTGGCTCG	ATACTGACGC	TGAGGTGCGA	700
	AAGCGTGGGG	AGCAAACAGG	ATTAGATACC	CTGGTAGTCC	ACGCCGTAAA	750
40	CGATGAATGC	CAGACGTCGG	CAAGCATGCT	TGTCGGTGTC	ACACCTAACG	800
	GATTAAAGCAT	TCCGCCTGGG	GAGTACGGTC	GCAAGATTAA	AACTCAAAGG	850
45	AATTGACGGG	GGCCCGCACA	AGCGGTGGAG	CATGTGGTTT	AATTCGAAGC	900
	AACGCGCAGA	ACCTTACCAA	CCCTTGACAT	GGCAGGACCG	CTGGAGAGAT	950
	TCAGCTTTCT	CGTAAGAGAC	CTGCACACAG	GTGCTGCATG	GCTGTCGTCA	1000
50	GCTCGTGTCG	TGAGATGTTC	GGTTAAGTCC	GGCAACGAGC	GCAACCCACG	1050

55

TCCCTAGTTG CCAGCAATTC AGTTGGGAAC TCTATGGAAA CTGCCGATGA 1100
 TAAGTCGGAG GAAGGTGTGG ATGACGTCAA GTCCTCATGG GCCTTACGGG 1150
 5 TTGGGCTACA CACGTGCTAC AATGGTGGTG ACAGTGGGTT AATCCCCAAA 1200
 AGCCATCTCA GTTCGGATTG TCCTCTGCAA CTCGAGGGCA TGAAGTTGGA 1250
 10 ATCGCTAGTA ATCGCGGAAC AGCATGCCGC GGTGAATACG TTCCCGGGCC 1300
 TTGTACACAC CGCCCGTCAC ACCATGGGAG TTGGTTCTAC CCGACGACGN 1350
 TGCCTAACC TTCGGGGGGC AGGCGGCCAC GGTAGGATCA GCGACTGGGG 1400
 15 TGAAGTCGTA ACAAGGTAGC CGTAGGGGAA CCTGCGGCTG GATCACCTCC 1450
 TT 1452

20

Claims

1. Bacteria belonging to a new genus and having the following properties:

25

30

35

40

45

(1)	Morphology	polymorphic rod
(2)	Motility	motile
(3)	Flagella	peritricous
(4)	Spore formation	none
(5)	Gram stain	negative
(6)	Production of pigment	positive (water insoluble)
(7)	Oxidase	positive
(8)	Catalase	positive
(9)	Behavior toward oxygen	aerobic
(10)	Fermentation of glucose	negative
(11)	Production of 3-ketolactose	negative
(12)	Quinone type	Q-10
(13)	GC content of intracellular DNA	64 - 69 molar%
(14)	Formation of slime	
	Glucose	negative
	Sucrose	negative
(15)	Presence of sphingolipid	negative
(16)	Presence of bacteriochlorophyll	negative

2. A bacterium to claim 1, wherein the bacterium is the strain E-396 (FERM BP-4283) or H-581-1 (FERM BP-4671).

50

3. A process for production of carotenoid pigment selected from the group consisting of astaxanthin, adonixanthin, β -carotene, echinenone, canthaxanthin and zeaxanthin, comprising the steps of:

culturing a bacterium capable of producing at least one of said carotenoid pigments, and recovering an individual carotenoid pigment or a mixture of the carotenoid pigments, wherein the bacterium has the following properties:

55

EP 0 635 576 A1

(1)	Morphology	polymorphic rod
(2)	Motility	motile
(3)	Flagella	peritricous
(4)	Spore formation	none
(5)	Gram stain	negative
(6)	Production of pigment	positive (water insoluble)
(7)	Oxidase	positive
(8)	Catalase	positive
(9)	Behavior toward oxygen	aerobic
(10)	Fermentation of glucose	negative
(11)	Production of 3-ketolactose	negative
(12)	Quinone type	Q-10
(13)	GC content of intracellular DNA	64 - 69 molar%
(14)	Formation of slime	
	Glucose	negative
	Sucrose	negative
(15)	Presence of sphingolipid	negative
(16)	Presence of bacteriochlorophyll	negative

4. A bacterium to claim 3, wherein the bacterium is the strain E-396 (FERM BP-4283) or H-581-1 (FERM BP-4671).

Fig.1




Fig.2




Fig.3




Fig.4




Fig.5





Fig.6



α -2, α -3, α -4, represent subclasses of Proteobacteria.
Phylogenetic Tree of E-396 and Close Species



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 94111525.5
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 6)
A	<u>AT - B - 304 414</u> (F. HOFFMANN-LA ROCHE & CO.) * Claim 1 * --	3	C 12 P 25/00 C 12 N 1/20
A	<u>WO - A - 91/03 571</u> (APPLIED FOOD BIOTECHNOLOGY) * Abstract * --	3	
A	<u>DE - A - 2 436 189</u> (SOCIETE DES PRODUITS NESTLE) * Claim 1 * --	3	
A	<u>DE - A - 2 436 188</u> (SOCIETE DES PRODUITS NESTLE) * Claim 1 * --	3	
D,A	APPLIED AND ENVIROMENTAL MICROBIOLOGY, vol. 55, no. 10, October 1989, Washington DC H.J. NELIS et al. "Reinves- tigation of Brevibacterium sp. Strain KY-4313 as a Source of Canthaxanthin" pages 2505-2510 * Totality * --	3	TECHNICAL FIELDS SEARCHED (Int. Cl. 6) C 12 P C 12 N
D,A	THE JOURNAL OF GENERAL AND APPLIED MICROBIOLOGY, vol. 15, no. 1, 1969, Tokyo H. IIZUKA et al. "Micro- biological Studies on Petroleum and Natural Gas; X. Carotenoid Pigments of Hydrocarbon-utilizing Bacteria" pages 127-134 * Totality * ----	3	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 06-10-1994	Examiner WOLF
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			